

**COLORADO RIVER RECOVERY PROGRAM  
FY 2008 ANNUAL PROJECT REPORT**

**RECOVERY PROGRAM  
PROJECT NUMBER: RZ-RECR**

I. Project Title: Razorback emigration from the Stirrup floodplain (RM 275.7)

II. Principal Investigator(s):

Trina Hedrick/Leisa Monroe  
Utah Division of Wildlife Resources  
Northeast Region  
152 East 100 North  
Vernal, Utah 84078  
Phone: (435) 781-9453 fax: (435) 789-8343  
E-mail: [trinahedrick@utah.gov](mailto:trinahedrick@utah.gov)  
[leisamonroe@utah.gov](mailto:leisamonroe@utah.gov)

III. Project Summary:

Floodplain wetlands are presumed to be important rearing habitat for razorback sucker (*Xyrauchen texanus*) (Wydoski and Wick 1998; Muth et al. 1998; Lentsch et al. 1996; Modde 1996; Tyus and Karp 1990). Reproduction by razorback suckers occurs on the ascending limb of the spring hydrograph allowing enough time between hatching and swim up for larvae to enter the system when highly productive floodplain habitats are accessible (Muth et al. 1998). This seasonal timing of razorback sucker reproduction indicates possible adaptation for using floodplain habitats for rearing purposes (Muth et al. 1998). It is currently unclear, however, how long young razorback sucker tend to stay in the floodplain before moving back out into the river.

The Green River Floodplain Management Plan (2003) identifies the Stirrup floodplain as a high priority habitat for recovery of the endangered razorback sucker, bonytail (*Gila elegans*), and Colorado pikeminnow (*Ptychocheilus lucius*). The natural levee surrounding the Stirrup was breached at the downstream end in March 1997 in an effort to increase the frequency of connectivity of the floodplain to the river. The floodplain now connects at around 14,000 cfs and can fill to approximately 20 acres during spring peak flows (Birchell and Christopherson 2004).

Though it is not extremely large, it is one of the few floodplain habitats in the middle Green River that retains enough water and overall depth to over-winter fish, thus making it ideal when maintaining razorback sucker over multiple years. Because of its potential to overwinter fish and because it only has one breach, this site was chosen for a study to research the timing of razorback sucker emigration from highly productive floodplain habitats to the river. Surplus razorback sucker were identified from normal operations at the Ouray National Fish Hatchery and were held at the hatchery site until they were stocked into the Stirrup floodplain: age-2 fish in June 2007 and age-1 fish in October 2007. These fish were PIT tagged for individual identification, stocked, and then monitored in the fall and spring. In May 2008, during spring peak flows, a stationary PIT tag reader was set up to monitor tagged fish movement into and out of the floodplain.

IV. Study Schedule: Initial year - FY - 2007 Final year - FY 2010 (final year was originally intended to be FY 2009)

V. Relationship to RIPRAP:

#### GENERAL RECOVERY PROGRAM SUPPORT ACTION PLAN

II. Restore habitat (habitat development and maintenance)

II.A. Restore flooded bottomland habitats

II.A.1. Conduct inventory of flooded bottomlands habitat for potential restoration

#### GREEN RIVER ACTION PLAN: MAINSTEM

II. Restore habitat (habitat development and maintenance)

II.A. Restore and manage flooded bottomland habitat

II.A.1. Conduct site restoration

II.A.2. Acquire interest in high-priority flooded bottomland habitats between Ouray NWR and Jensen to benefit endangered fish

II.A.2.a. Identify and evaluate sites

IV. Manage genetic integrity and augment or restore populations (stocking endangered fishes)

VI. Accomplishment of FY 2008 Tasks and Deliverables, Discussion of Initial Findings and Shortcomings:

**Task 1.** Pump water from the river into the Stirrup floodplain. This includes preparation of compliance documents for both the BLM and Utah Division of Water Rights. This may also be done again between tasks 3 and 4. Spring and summer (potential to require pumping at other times of year as well)

The U.S. Fish and Wildlife Service (Program Director's Office and Ecological Services) completed the EA to do the work on Bureau of Land Management land in August 2007. The Division obtained a temporary water right and began pumping on 20 June 2007 with two Honda WT40X, 4" trash pumps; however, after a number of weeks pumping with the 4" pumps, it became clear that we were not going to increase the depth of the floodplain with these small pumps.

Pumping with a 6" trash pump began on 15 October and ended on 7 November 2007. Depth at this time was 4.5' and was determined to be sufficient to maintain fish overwinter.

Pumping occurred again in the early spring before connection with the river. Pumping was done using the 4" trash pumps and was only to provide an influx of riverine water that had higher DO levels than what was present in the Stirrup after ice-off. Pumping continued for two weeks.

Pumping after spring 2008 river connection was not necessary as water quality and overall depth remained adequate (see Task 3).

### **Task 2. Stock razorback sucker in the Stirrup floodplain**

The Ouray National Fish Hatchery stocked age-1 and age-2 fish in the Stirrup in 2007. These were the fish we attempted to detect with the stationary PIT tag reader during connection in 2008.

After connection in 2008, hatchery personnel stocked an additional 1,047 PIT-tagged, age-1 razorback sucker (2007 cohort; avg TL = 203mm); and 952 PIT-tagged, age-2 fish (2006 cohort; avg TL = 309). In October 2008, approximately 1000 PIT-tagged age-0 (2008 cohort) were also stocked. Note that the 2008 stocking events occurred on top of an undetermined number of survivors from the 2007 stockings that remained in the Stirrup after the high flows of 2008.

### **Task 3. Monitor water quality and species assemblage in Stirrup floodplain**

While the floodplain did not freeze all the way through this winter, low dissolved oxygen was a problem due to a lack of sunlight getting through to algae in the floodplain. Testing on 31 January resulted in 11" of ice and 7" of snow on top of the ice making the floodplain nearly void of sunlight. Dissolved oxygen readings taken on 14 February were less than 1.0 mg/L.

We sampled the Stirrup with fyke nets on 9, 10 April. While we did not capture any razorback suckers, we did see a number of dead razorback suckers and carp along the northeast shoreline. There was a substantial fish kill due to low dissolved oxygen overwinter. The depth in the floodplain made sampling efficiently with fyke nets difficult; we therefore sampled with trammel nets, electrofishing boats and a combination of the two to determine whether the fish kill was complete or not. These methods were unsuccessful and revealed few bullhead, catfish, and fathead minnows in the floodplain.

After spring connection, the Stirrup was completely full and held water quite well over the summer. Dissolved oxygen (DO) was measured on 5 August 2008. The measurement recorded was 12.29 mg/L at 17:14. DO was again measured on 28 August 2008. The measurement was taken at 15:20. Temperature was 21.42°C and DO was 10.72 mg/L. DO tends to be highest around 23:00 in the Stirrup. These measurements are therefore more than adequate to maintain razorback sucker overwinter if the winter is not as harsh.

### **Task 4. Research stationary PIT tag readers and determine the appropriate set up for the Stirrup floodplain**

Crews were sent to the San Rafael drainage and the Provo River drainage to see other stationary PIT tag readers being set up. Through this and speaking with representatives from Biomark, we decided to use the FS1001M Reader (MUX), which could monitor up to six antennas at once. This first year, we decided to begin with only one antenna; however, in the future, we will build more. Crews sent out to the Stirrup breach decided that a 10' x 3' antenna would adequately

cover the breach, although once the breach had filled, a larger antenna would have also been useful.

**Task 5.** Set up stationary PIT tag reader during spring peak flows.

The Stirrup first connected to the floodplain on 11 May, 2008. This connection did not last long and since we were experiencing problems with the reader (see below), four trammel nets were set within the breach to prevent fish from moving into or out of the floodplain. These trammel nets were successful at restricting movement during the entire first connection, which only lasted one day. Four carp were captured attempting to move out of the floodplain and one channel catfish was captured trying to move into the floodplain.

By 20 May, the beginning of the second connection, the reader was set up and functioning adequately. The first incident of noise was worked out (see description of problems with the plug or connector) and the reader was sampling the breach for at least the first 48 hours of connection as displayed by the capture of a test tag. On 22 May, we realized that we were going to have battery issues and were going to have to replace the batteries every day as the solar panel was not charging the batteries as it should (see description below). On 23 May, we moved the reader to higher ground as we underestimated the magnitude of flows down the river and the MUX was now in danger of being swamped. When we got the MUX to higher ground and had everything running again, a serious level of noise (93%, the maximum noise reading) inhibited our ability to read tags until 1 June (see below for this explanation). We finally figured out the cause of this noise and had the reader running with no problems through 7 June. During this time, the MUX picked up 6 tags. On 8 June, we had low current readings from the antenna; the description of how this was fixed is included below. The antenna was back out and ready to go on 9 June. The MUX picked up two more tags on 10, 11 June. We had one more noise issue the evening of the 11<sup>th</sup> (see Dynamic Tuning description below) that we were able to correct, though we lost one more night due to this issue). The antenna was taken down on 14 June (Jensen gauge reading was 14,300 cfs and coming down) due to only a small amount of very clear water in the breach. No more tags were picked up, likely because the water in the breach was quite shallow and clear beginning on 11 June. Below is a list of difficulties experienced during 2008 high flows:

- Noise issues –
  - The antenna connects to the MUX with a large cable that has one black wire, one red wire, and a metal sheath around these two wires that is formed into a wire for the ground. Connector pins are soldered onto these wires, which are then inserted into a plastic connector or plug that tightens down onto one of the MUX antenna ports. It was very difficult to adequately solder and secure the pins, insert them into the plug or connector so each pin extended out of the plug the same distance, and ensure a solid connection without getting a high noise reading. We ended up scrapping the plastic connector and inserting the pins into the MUX antenna port without a plastic connector. Because the MUX (and therefore the end of the antenna cable) is secured within a metal enclosure, there was no real need for a connector. By scrapping the connector, we were able to reduce the noise (though it was not obvious, the wires must have been touching somewhere within this

connector) and ensure a solid connection. This issue was resolved before connection occurred.

- Dynamic Tuning also created noise high noise levels. There are two different ways to tune the antenna; one requires dynamic tuning to be turned on and the other requires dynamic tuning to be turned off. Because the dynamic tuning feature is always on unless the user specifically turns it off, every time we went into the manual tune option (with a hyperlink between the MUX and a laptop), the dynamic tune option would turn back on and increase the noise level. It was actually quite some time before we realized this was the reason for our high noise levels. Before pinpointing this as the reason, we tried moving the antenna and removing every piece of metal near the antenna. We also tried using the solar panels without a regulator in case that was increasing the noise level. We replaced batteries and checked every connection. We inserted the antenna cable into a different port. We re-soldered the connector pins to the antenna cable wire a number of times as well. Throughout this entire ordeal, we were in close contact with the Division's Springville office and the USGS fisheries station in Klamath as both shops have used this technology and are familiar with many of the problems that can occur. They recommended a number of things, but it wasn't until someone suggested ensuring the Virtual Test Tag was off that we tried turning off the Dynamic Tune, which immediately resulted in a decrease in the noise level.
- Drained batteries –
  - We used two deep-cycle marine batteries to power the MUX. These batteries will continue to power the MUX with > 20V; however, even in direct sunlight, the solar panel was only re-charging the batteries for about 1.5 days. We were continually replacing batteries to keep the MUX going. By doing this, we didn't lose much sampling time to dead batteries; however, in the future, we will purchase marine deep cycle batteries with greater amp-hours in order to increase the time between battery replacements. The Division's Springville office ran their MUX on the same two batteries for nearly three months without having to replace them. We should therefore not have to replace the batteries at all for this project.
- Reduced antenna current –
  - To pick up a tag, the antenna current must be between 3.0 and 6.0 amps. After we figured out the noise issue and realized we were going to have to replace the batteries every day, the next problem was a low antenna current warning. We removed the antenna from the breach, brought it back to the shop, and opened the end cap. Water had been seeping in very slowly, but it was wet enough inside to inhibit the current output. We let the antenna dry out, resealed it with glue (we had used thread seal before) and replaced it in the breach.

#### **Task 6.** Download PIT tag data and monitor PIT tag array

The list of fish picked up with the MUX is as follows (directionality is assumed):

A razorback sucker (3D9.1BF18DFC0A) entered the breach on 2 June at 14:21 and left again at 16:01. This fish came back on 3 June and was picked up at 03:24, 03:45, and at 10:48. This fish was 425mm when we originally tagged it above the Jensen Bridge in 2007 (near river mile (RM) 301.8). We did not detect an old tag in this fish.

A bonytail (3D9.257C669F5B) entered the breach at 02:17 on 3 June. This was a 206mm fish stocked by Wahweap hatchery in 2007 at Red Wash (RM 298.1) in the middle Green River.

A Colorado pikeminnow (3D9.1BF18E65F7) entered the breach at 12:51 on 3 June. He subsequently left at 14:15. This fish was again detected on 4 June at 10:52, 11:33, 14:19, and 15:47. He was originally tagged at 460mm this year during the Colorado pikeminnow abundance estimates at river mile 275.6.

A Colorado pikeminnow (3D9.1BF234E6B0) entered the breach at 00:11 on 4 June. He was 517mm at the time of his original tagging in 2007 just below the Stirrup breach. No old PIT tag was detected.

A razorback sucker (3D9.1BF18CF5F7) was detected in the breach on 5 June at 04:02, 05:53, 06:22, 11:56, and 15:38. This fish was originally tagged in 2005 by Ron Brunson during northern pike removal by Razorback Bar (RM 311.0). He was 445mm at the time of tagging and has not been captured since then.

A Colorado pikeminnow (3D9.257C66D4FE) was captured going into the floodplain at 00:17 on 7 June. This fish was originally tagged by Paul Badame's crew last May at RM 74.5. It was 435mm at the time of tagging.

A roundtail chub (3D9.1BF1FF6409) was captured heading into the floodplain sometime between 18:00 on 10 June and 09:45 on 11 June (the batteries had died previous to this and we had forgotten to reset the date/time feature). This fish was originally tagged this year during the Colorado pikeminnow abundance estimates at RM 275.1. It was 378mm at the time of tagging.

A razorback sucker (3D9.257C698F18) was captured leaving the floodplain on 11 June at 22:23. This was an age-1 fish stocked into the Stirrup in October 2007 specifically for this study. This was the only fish stocked into the floodplain that was detected moving out during high flows.

#### **Task 7. Summarize results/findings**

Annual report submitted November 2008.

#### **VII. Recommendations:**

- Use 4” trash pumps only to “top off” the floodplain; use 6” trash pumps when trying to add substantial depth to and improve water quality of the floodplain.
- Be flexible on the sampling methods chosen for the Stirrup. For example, before connection with the river, electrofishing may not be effective due to low conductivity; however, this may be a more effective method than fyke netting or trammel netting after connection occurs.
- Build two more antennas for the breach to be able to determine directionality and probability of detection for each antenna.
- Continue to use deep cycle marine batteries, but purchase batteries with more amp hours than those that were used this year.
- Explore different means of securing the antenna in the breach. Metal objects can create noise in the antenna. The bottom of the breach is also quite sandy, which made securing the antenna difficult with other means. However, due to read range, we need to have the antenna upright so we need to have more and better options available to next year.
- Have the MUX in the Stirrup breach one week before an expected connection in order to work out the few remaining difficulties left over from sampling in 2008 (batteries, etc.).

VIII. Project Status:

Ongoing

IX. FY 2008 Budget Status

- A. Funds Provided: \$36,126
- B. Funds Expended: \$36,126
- C. Difference: \$0
- D. Percent of the FY 2008 work completed, and projected costs to complete: 100%
- E. Recovery Program funds spent for publication charges: \$0

XI. Signed: Trina Hedrick 10/31/08  
Principal Investigator Date

XII. Literature Cited

Birchell, G.J. and K. Christopherson. 2004. Survival, growth, and recruitment of larval and juveniles razorback sucker (*Xyrauchen texanus*) introduced into floodplain depressions of the Green River, Utah. Utah Division of Wildlife Resources, publication no. 04-15, Salt Lake City, Utah.

Christopherson, K. Native Aquatics Project Leader, Utah Division of Wildlife Resources, Northeastern Region, 1998-2006.

Valdez, R.A. and P. Nelson. 2003. Green River Subbasin Floodplain Management Plan. Biology Committee Review Draft, R.A. Valdez & Associates, Inc., Upper Colorado River Endangered Fish Recovery Program, Denver, CO.

Lentsch, L., T. Crowl, P. Nelson, and T. Modde. 1996. Levee removal strategic plan. Utah Division of Wildlife Resources, Salt Lake City, UT. 21 pp.

Modde, T. 1996. Juvenile razorback sucker (*Xyrauchen texanus*) in a managed wetland adjacent of the Green River. Great Basin Naturalist 56:375-376.6

Muth, R.T., G.B. Haines, S.M. Meismer, E.J. Wick, T.E. Chart, D.E. Snyder, and J.M. Bundy. 1998. Reproduction and early life history of razorback sucker in the Green River, Utah and Colorado, 1992 – 1996. Final Report submitted to the Recovery Implementation Program for Endangered Fish Species in the Upper Colorado River Basin. U.S. Fish and Wildlife Service, Denver, CO. 62 pp.

Tyus, H.M. and C.A. Karp. 1990. Spawning and movements of razorback sucker, *Xyrauchen texanus*, in the Green River basin of Colorado and Utah. Southwestern Naturalist 35:427-433.

Wydoski, R.S. and E.J. Wick. 1998. Ecological value of floodplain habitats to razorback suckers in the Upper Colorado River Basin. Final Report submitted to the Recovery Implementation Program for Endangered Fish Species in the Upper Colorado River Basin. U.S. Fish and Wildlife Service, Denver, SO. 55 pp.