

COLORADO RIVER RECOVERY PROGRAM  
FY 2016 ANNUAL PROJECT REPORT

RECOVERY PROGRAM  
PROJECT NUMBER: FR-115

- I. Project Title: Monitoring effects of Flaming Gorge Dam releases on the Lodore and Whirlpool Canyon fish communities
- II. Bureau of Reclamation Agreement Number(s): R14AP00001  
Project/Grant Period: Start date (Mo/Day/Yr): 1 Oct. 2014  
End date: (Mo/Day/Yr): 30 Sept. 2018  
Reporting period end date: 30 Sept. 2016  
Is this the final report? Yes \_\_\_\_\_ No X
- III. Principal Investigator(s): Lead Agency: Larval Fish Laboratory, Department of Fish, Wildlife, and Conservation Biology, Colorado State University; Bureau of Reclamation; U.S. Fish and Wildlife Service

Kevin R. Bestgen, E. Kluender and K. Zelasko  
Larval Fish Laboratory  
Dept. of Fish, Wildlife, and Conservation Biology  
Colorado State University  
Ft Collins, Colorado 80523  
970-491-1848/ fax 970-491-5091  
kbestgen@colostate.edu

Tildon Jones  
Vernal Colorado River Fish Project  
U.S Fish and Wildlife Service  
1380 S. 2350 W.  
Vernal, Utah 84078  
435-789-0354/ fax 435-789-4805

- IV. Abstract: The primary purpose of this study is to determine the cumulative effect that flow and temperature regimes downstream of Flaming Gorge Dam have had on the fish community of the Green River in Lodore and Whirlpool canyons and to recommend how to monitor effects into the future. A secondary purpose is to determine the distribution and abundance of the humpback chub population in Whirlpool Canyon to serve as the basis for future monitoring efforts. Finally, a third purpose is to remove nonnative fishes present in the study reach. Future monitoring will be needed to evaluate the contribution of the Whirlpool Canyon population of humpback chub to the overall recovery of the species, and to monitor response of the fish community to removal of smallmouth bass, particularly in the Whirlpool Canyon reach of the Green River. A portion of nonnative fish removal work is devoted to better understanding the reproductive ecology of smallmouth bass in the Green River study area. The collection of young-of-year smallmouth bass, and analysis of otolith microstructure, will allow determination of hatching dates of bass relative to streamflow and water temperature patterns. This

information may be useful to understand if flow releases from Flaming Gorge Dam may be useful to disadvantage smallmouth bass in the study area. Northern pike are also a species targeted in removal efforts and occur mainly in upstream Browns Park. Information gathered will be used to evaluate the relative benefit of flow and temperature regimes from Flaming Gorge Dam to native and endangered fishes in the Green River compared to effects on distribution and abundance of non-native fishes.

V. Study Schedule: Initial Year: 2002; Final Year: Not determined

VI. Relationship to RIPRAP:

Green River Action Plan: Mainstem.

II.D. Evaluate and revise as needed, flow regimes to benefit endangered fish populations.

VII. Accomplishment of FY 2016 Tasks and Deliverables, Discussion of Initial Findings and Shortcomings:

Task 1: Thermographs

We assisted with temperature data collection in Lodore and Whirlpool canyons by removing thermographs from the river in July and September<sup>1</sup>, resetting new ones in their place, and establishing/checking new locations in both lower Lodore Canyon near the existing site, and in lower Mitten Park on river right in 2014<sup>2</sup>; that monitoring continues through 2016. Over the past decade, the thermal regime of the Green River upstream of the Yampa River has varied. The section experienced a relatively warm regime from 2002-2007 (2005 was cool) when water temperatures routinely exceeded 20°C at the Gates of Lodore, and cool thermal regimes in much of the 2008-2011 period, when water temperatures exceeded 20°C for just a few days each summer (Figure 1). Water temperatures were warmer in 2012 and 2013 but cooler in 2014-2016, when mean daily water temperature did not or rarely exceeded 20°C at the Gates of Lodore (gauge 404417108524900).

In 2016, Green River water temperatures at the Gates of Lodore were increasing in May consistent with warming air temperatures to maxima near 20C, but during high flow releases (up to 9100 cfs) from late May until the end of June, declined to less than 12°C (Figure 1B). Comparison of water temperatures from releases at Flaming Gorge Dam to the Gates of Lodore (Figure 1B & 1C) showed little warming downstream in those high flow periods. However, water temperature recovered downstream after high flows ceased. Highest water temperatures in 2016 were in mid-July but daytime maxima exceeded 20°C for several days in August as well.

---

<sup>1</sup> Thermograph data will be provided by U.S. Fish and Wildlife Service, Lakewood, and by Utah State University, at up to 10 other localities in the Green River.

<sup>2</sup> That Mitten Park location is across the river from the long-term site, but had to be relocated due to sedimentation.

## Task 2: Sample main channel fish community (large-bodied fishes).

We completed two electrofishing trips through the study area in 2016, as prescribed in the study proposal. We reported on data collected through 2006 in a summary report. Data entry and verification is complete for 2007-2015; 2016 data is preliminary but is included in this report. We are in the process of summarizing data for an updated synthesis report, which will include electrofishing data through 2016.

A total of 11 native species, 20 nonnative species, and several hybrids have been collected in the study area by all sampling gears in the period 2002-2016 (Table 1, Figures 2 and 3). White crappie is a new species, collected with seines in Whirlpool Canyon. The most species are captured by electrofishing, followed by seining, and trammel netting. In 2016, the most abundant species in the Lodore Canyon study area captured by electrofishing were native flannelmouth and bluehead suckers (40% in 2016, 56% in 2015); native fishes were 54% (61% in 2015) of all fishes captured. Brown trout, channel catfish, white sucker, common carp, and smallmouth bass, in descending order, were the most abundant nonnative fishes captured in the study area.

Native fishes were more abundant in downstream Whirlpool Canyon, where 70% (71% in 2015) of all fishes collected by electrofishing were native, which was substantially higher than in upstream Lodore Canyon. A main difference was higher abundance of bluehead sucker in Whirlpool Canyon. Conversely, coolwater mountain whitefish was more abundant in Lodore Canyon than Whirlpool Canyon. Overall, Whirlpool Canyon sampling captured only about 50% of the fish captured in Lodore Canyon, which was due in part to differences in sampling effort and reduced abundance of brown trout downstream. Among nonnative fishes, coolwater brown trout were more abundant in upstream Lodore Canyon and warmwater smallmouth bass and channel catfish were more abundant downstream in Whirlpool Canyon.

Abundance patterns of fishes in the study area that were captured in electrofishing samples beginning in 1994-1996 reflected a dynamic community for both nonnative and native fishes (Figures 2 and 3). For example, nonnative brown trout was abundant through time in Lodore Canyon and apparently increased in 2012, but declined in abundance in 2013-2106 to relatively low levels. Brown trout abundance increased slightly in Whirlpool Canyon in 2012-2014 but was uncommon in 2015 and 2016. Only in 2008 was that species abundant there due to large numbers captured near Jones Creek, a location sampled only sporadically because of presence of fisherpersons. Channel catfish reached a peak in abundance in both Lodore and Whirlpool canyons in the 2005-2008 period but have declined steadily since, although abundance in 2015 in Whirlpool Canyon increased. Note the 1994-1996 abundance as the lowest in the period of record in Lodore Canyon. Common carp abundance appears to be declining in both reaches over time, especially in Whirlpool Canyon, although 2015-2016 abundance in each reach increased. Northern pike abundance is low overall but highest in Lodore Canyon and declining over time. The spike in 2016 abundance in Lodore Canyon was from only five individuals. Northern pike abundance in Whirlpool Canyon is low and none were captured in 2013-2016, likely a byproduct of high removal in the upstream Yampa River

which reduces movement into downstream reaches. Browns Park pike removal efforts in 2011-2016 are discussed more extensively below.

Smallmouth bass abundance in Lodore and Whirlpool canyons showed slightly different patterns. In Whirlpool Canyon, bass abundance peaked in 2007, and declined through 2011. Bass abundance then increased in 2012 and 2013, but declined in 2014-2016. That 2013 increase may be the result of warm water temperatures in 2012, leading to successful spawning that produced many age-1 fish in 2013. While earlier declines may reflect substantially increased removal effort beginning in 2007 and lack of high recruitment through 2011, bass respond positively to warmer temperatures in some years. Whirlpool Canyon bass abundance declines may be due to continued and intensive removal there. Smallmouth bass abundance in Lodore Canyon peaked in 2009 and 2010, declined through 2012 but then increased slightly in 2013, and then continued lower through 2016. White sucker abundance increased in Lodore Canyon through 2009 but then declined through 2015, with a slight increase in 2016. With the exception of 2008, white suckers in Whirlpool Canyon were relatively uncommon through 2016.

Native bluehead sucker are abundant in both reaches, but abundance is higher in Whirlpool than Lodore Canyon. Although abundance declined through 2012, abundance increased in 2013 and has remained higher since then. Bluehead sucker abundance in Lodore Canyon stabilized or increased slightly from 2009-2013 after a decline in abundance from the mid-1990s, declined again in 2014 and 2015, but increased in 2016. Colorado pikeminnow abundance has been variable and relatively low over the study period, increased in 2010 and 2012 in each reach, low overall in 2015, but increased in 2016 (n = 6 fish). The finding of pikeminnow in upstream Vermillion Creek in 2011-2016 is discussed below. Flannelmouth sucker, the most common large-bodied native fish in the study area, was been declining over time in Lodore Canyon, but increased in 2015 and 2016 to average or above average levels. Flannelmouth sucker was very abundant in 2008 in Whirlpool Canyon but declined over time until 2011 after which abundance increased steadily through 2016. Mountain whitefish abundance increased rather dramatically in the study area in some years such as 2009-2010, 2012, and 2016 but were uncommon in 2013-2015. Decreased salmonid abundances in both reaches in 2013-2015 may be a response to warmer water temperatures.

Abundance of all chub species declined rather dramatically, especially since 2002-2004, patterns also reflected by trammel net sampling. Capture of a few chubs in 2013 electrofishing in Lodore Canyon increased abundance there temporarily, but 2014 abundance was reduced in both reaches, none were captured in either Lodore or Whirlpool Canyon reaches in 2015, and only one was captured in 2016, in Whirlpool Canyon.

A dedicated trammel net sampling trip occurred on 27 September to 30 September, where three locations in Whirlpool Canyon were sampled. We captured a total of one roundtail chub (not a recapture) and 3 individual bonytail (all recaptured). The roundtail chub was 351 mm TL, and bonytail were 250 - 403 mm TL.

Beginning in 2005, northern pike were detected nearly every year in Browns Park. Initially, only smaller pike were captured in seine samples taken in backwaters in July and September at Swinging Bridge near the Colorado-Utah state line. Their continued presence motivated additional floodplain sampling to attempt to remove as many adults as possible. Thus, beginning in 2011 and continuing through 2016, we increased sampling effort for removal of northern pike in the Browns Park reach of the Green River. During the high flow year 2011 in May and June many large pike, some exceeding 900 mm TL (n=22, 11-39 inches, 271-984 mm) were captured and removed; all were from a relatively restricted river reach about 1-mile long. During that sampling, two adult Colorado pikeminnow *Ptychocheilus lucius* were captured, one by angling and another in a trap net. This was unusual because pikeminnow are thought rare in the 56-mile long reach between Flaming Gorge Dam and the upstream end of Lodore Canyon which includes the Swinging Bridge area. In 2012, we captured additional northern pike in the Browns Park reach during a relatively brief period of high flows. Interestingly, only three young-of-year were captured in summer and autumn, which was perhaps a function of the brief period of floodplain inundation. No young of year northern pike were captured in 2013 in Browns Park, again likely a result of relatively short-term floodplain inundation. This is in spite of capture of ripe northern pike in late May and early June, indicating reproduction was likely taking place. More details on pike abundance and spawning patterns were presented at the non-native fish workshop in December 2013.

In 2014, continued backwater and floodplain habitat sampling occurred with trammel and fyke nets from 26 May to 20 June. A total of 12 adult northern pike was captured and ranged from 626-794 mm TL (1814-3657 g). Most were captured in Beaver Creek (n = 7), but northern pike were also captured at or near Hog Lake (n = 2) or in Vermillion Creek (n = 3). The Vermillion Creek pike were the first ones captured there in any sampling period. In spite of warm water temperatures earlier in the season, many pike were still in reproductive condition in late May to mid-June. Of the ten fish whose sex was determined, six were females.

We captured a total of 13 age-0 northern pike in 13 July 2014 seine sampling in Browns Park at the mouth of Beaver Creek, ranging in size from 44 to 61 mm TL. An additional seven age-0 northern pike were captured in seine sampling on 14 September at Beaver Creek (n = 6) and Hog Lake (n=1), ranging in size from 174 to 218 mm TL.

Northern pike sampling and removal in 2015 began on 12 May and continued through 9 June. Due to extensive habitat availability and high initial fish catch, sampling was focused primarily in the State Line and Beaver Creek reach; all other accessible backwaters and tributaries in the Refuge were sampled opportunistically during this period, however. Nine adult northern pike (289 – 776 mm, 260 – 2800 g) were captured. As in 2014, most were captured in Beaver Creek (n = 4), but the State Line backwater (n = 2), a backwater near Spitzie Bottom (n = 2), and Vermillion Creek (n = 1) produced pike. All adult northern pike were male. In addition to trammel netting for adults, intensive nighttime light-trapping for age-0 pike was conducted for the entire sampling

period. As with trammel netting, effort focused on the State Line backwater to Beaver Creek reach, but some efforts were made in other backwaters.

Light trapping in 2015 produced one age-0 northern pike (30 mm TL) in Beaver Creek on 23 May. A single age-0 northern pike was collected in seine sampling in Browns Park on 19 July. This individual was 126 mm and was captured near the boat ramp at Crook Campground.

Sampling in 2016 in Browns Park National Wildlife Refuge occurred between 11 May and 18 June 2015. Sampling was carried out by a crew of two to six individuals. Trammel nets measuring 15 m x 2 m were the primary gear used for adult fish, with some supplementary angling. A total of 54 net sets were made, totaling 353.67 net hours. Two boat electrofishing passes were conducted; the first covered the left bank between river miles 260.2 (Swinging Bridge Boat Ramp) and 252.1 (Crook Campground Boat Ramp) and the second was conducted between river miles 252.1 and 246 and included the lower 400 meters of Vermillion Creek. Boat electrofishing effort totaled 3.44 hours. Light traps were deployed in backwaters at night to detect the presence of larval fish, northern pike being the primary target species. We focused light trap sampling efforts in the backwaters and main channel shoreline between the state line and the Swinging Bridge boat ramp. Up to five light traps were deployed most nights, and effort totaled 743.97 trap hours.

During the study, 869 individual adult fish were captured (605 by trammel nets, 265 by electrofishing, Appendix I). Of these, 333 were native species, hybrids of native species, salmonids, or unidentified, and were all released alive. The remaining 536 individuals were nonnative species and were euthanized. Light traps yielded a total of 48 fish; all were preserved and identified in the lab.

The discovery of Colorado pikeminnow upstream near Swinging Bridge in spring 2011 motivated additional Green River sampling in Browns Park National Wildlife Refuge on 21-22 June via boat electrofishing. Two crews sampled from Swinging Bridge downstream 14 miles to Vermillion Creek, a tributary to the Green River a short distance upstream of the boundary of Dinosaur National Monument. High flows benefitted sampling efforts because of increased access to productive habitats. One adult pikeminnow was captured in the Green River near Crook Campground. High Green River flows also allowed access to the lower end of Vermillion Creek, where an additional 8 adult pikeminnow (23-27 inches, 595-692 mm total length,) were captured; at least three more were observed but not captured. In all, only five pikeminnow had been previously tagged and two of those were tags with number series that have not been used for many years, which may indicate that those fish reside in that section of the Green River for substantial periods of the year. Vermillion Creek, a relatively small, turbid, and low flow system, was relatively warm at 72°F (22°C) compared to the Green River (48°F, 9°C); during these captures several individual pikeminnow were tuberculate (nearing reproductive readiness) likely because of the warm water. High reproductive condition for pikeminnow was not expected because of high, late, and cold flows and because pikeminnow in the downstream Yampa River did

not spawn until late July in 2011. Very high densities of suckers, both native flannelmouth sucker *Catostomus latipinnis*, non-native white sucker *Catostomus commersonii*, and their hybrids, were also captured. Findings also supported the importance of floodplain wetlands and flooded tributary mouths for enhancing condition of endangered fishes like Colorado pikeminnow. In spring 2012, we captured three additional pikeminnow at or in the mouth of Vermillion Creek, in spite of relatively low effort. Consecutive years of capture at that location suggest that the mouth of Vermillion Creek may be a concentration area for pikeminnow in spring.

In spring 2014, we captured six Colorado pikeminnow in or at the mouth of Vermillion Creek in fyke nets; an additional pikeminnow was captured at Beaver Creek. Captures were from 28 May to 11 June and each fish was captured only once. Colorado pikeminnow ranged in length from 502 – 756 mm TL (1162-4309 g) and six of the seven were previously tagged; one was tagged for the first time. One of those individuals that we captured (8 June) was also detected by a PIT tag detection array set in Vermillion Creek a few days earlier (4 June)<sup>3</sup>. Two additional individuals were also detected by arrays, but we did not capture those fish. Thus, a total of nine Colorado pikeminnow adults was captured in 2014 between 1-16 June and eight were from Vermillion Creek.

On 3 June, flows increased from Flaming Gorge Dam to 4,350 cfs, peaked on 7 June at 9,090 cfs and by 18 June flows were declining to or were below Flaming Gorge Dam maximum powerplant flows of 4,650 cfs or less. One pikeminnow was detected in the mouth of Vermillion Creek on 28 May at low base flows of 872 cfs. All pikeminnow except the first one were captured in or near Vermillion Creek during high flows (2-18 June) when the mouth of Vermillion Creek was inundated, and the Green River was cold (about 10°C). Vermillion Creek was warm and > 20°C for most of that high flow period.

In 2015, 17 individual Colorado pikeminnow (483–713 mm, 1300–3650 g) were captured in Browns Park. Of these, 12 were captured in Vermillion Creek within 100 m of the mouth over a span of five days (4 – 8 June). Four of the Vermillion Creek individuals were captured twice or more during this time. The remaining five Colorado pikeminnow were captured throughout the Refuge at State Line backwater (n = 1), Beaver Creek (n = 2), near Spitzie Bottom (n = 1), and in a backwater above Crook Campground (n = 1). Only three individuals were not tagged; these were given new PIT tags at capture. With the exception of one individual angled at Beaver Creek, all were captured with trammel nets. The Vermillion Creek individuals were captured after high-volume releases from Flaming Gorge had concluded, and water temperatures at the time of capture ranged from 13.9 to 24.4°C. Daytime main channel temperatures were not taken during this time but just downstream, minimum and maximum daily water temperatures were cooler at about 13-17C (Figure 1b).

Two electrofishing trips were conducted through Brown's Park in June to access backwaters and other habitats otherwise inaccessible for trammel netting. On 3 June, sampling was conducted with a raft-based Smith Root GPP electrofisher between the

---

<sup>3</sup> both a flat plate and a submersible antenna was deployed in the mouth of Vermillion Creek on 3 June (A. Webber, U.S. Fish and Wildlife Service, Vernal, Utah)

Swinging Bridge and Crook Campground boat ramps (river miles 260.3 to 252.1) and focused on backwaters and vegetated shoreline on the left bank. Sampling for this trip totaled 9177 seconds. On 18 June, sampling was conducted from the Crook Campground boat ramp to just below Vermillion Creek (river miles 252.1 to 246.0), focusing on backwaters and vegetated shoreline on both banks and on the lower 400 meters of Vermillion Creek. Sampling totaled 1398 seconds above Vermillion Creek, 1446 seconds in Vermillion Creek, and 368 seconds below.

In 2016, all 10 Colorado pikeminnow captured with electrofishing and trammel nets were captured in the lower 400 meters of Vermillion Creek during higher flow releases from Flaming Gorge Dam; one of these individuals was captured twice. During trammel netting in Vermillion Creek, water temperatures were between 17.0 and 20.9 degrees Celsius. Main channel Green River water temperatures were not taken on those days, but were measured between 8.1 and 9.9 degrees Celsius during June sampling. Two PIT tag antennas were deployed in lower Vermillion Creek to learn more about use of the tributary by Colorado pikeminnow and other tagged species of interest in the river. A total of 62 detections were made by the antennas, comprising 24 individuals. Of these fish, 20 were known Colorado pikeminnow and one was a known flannelmouth sucker. The species of the remaining three individuals are currently unknown. Five of the pikeminnow captured in Vermillion Creek (four by trammel nets and one by boat electrofishing) were also detected by the PIT tag antennas between 27 May and 17 June. Individuals detected by the PIT tag antennas were detected between 1 and 9 times, with a mean detection frequency of 2.7 times.

### Task 3: Sample small-bodied fish community.

A total of 232 seine samples were collected in the study area from middle Browns Park downstream to the lower end of Rainbow Park during summer and autumn 2016. We have begun identification of summer and autumn samples. We will update seine capture data when 2015 samples are identified.

An additional task in this scope of work was to analyze otoliths from age-0 smallmouth bass captured in the Green River study area. This work will assist with understanding smallmouth bass spawning periodicity to assist with disruption of reproduction of that species via flow releases from Flaming Gorge Dam. Studies in other parts of the range of smallmouth bass have shown that weather-related water temperature reductions or floods reduce their spawning success and number of offspring (See references in Bestgen and Hill 2016, smallmouth bass report). Reduced water temperatures often result in abandonment of spawning nests by the guarding male bass, after which developing eggs and just-hatched young are susceptible to predation and other mortality factors. Sampling in the Green River and other areas has shown that higher stream flow, often coupled with increased water turbidity, sweeps weak-swimming young bass away from nests or quiet near-shore habitat, and results in high mortality.

Portions of this work were reported at the January 2010-2016 Researchers Meetings. We

found that smallmouth bass in the Green River-Lodore Canyon study area first hatched well after spring peak releases declined and just slightly after (usually within one week) mean daily water temperatures regularly exceeded 16°C in the period 2003-2012 (Figure 4, note axis change for 2011 to accommodate the late spawning initiation). Hatching date distributions were very similar in 2008 and 2009, reflecting the similar flow and temperature regimes in those years. Hatching date and the extent of the reproductive season was much shorter in 2011 when high flows were relatively cool, and hatching did not begin in 2011 until after bass had *finished* hatching in nearly every other year. The 2012 water temperatures suggest a relatively late initiation of hatching as well, and that is supported by capture of very few and relatively small bass in Lodore Canyon in a late-July sampling trip. We have submitted and obtained approval of a report that synthesizes smallmouth bass hatching date distribution and growth data (Bestgen and Hill 2016).

Drift net sampling documented high downstream displacement of small-bodied smallmouth bass during high turbidity and flow events in 2004 and 2007. Such flow and turbidity events may have been responsible for low abundance of smallmouth bass < 100 mm TL in summer 2004, and subsequent low number of Age-1 smallmouth bass in 2005 (data in RIP annual rpts. Badame et al. synthesis report; discussion in Bestgen et al. 2006; 2007). Unfortunately, high flows in 2011 prevented sampling of the Green River because sampling crews were unable to cross the very high Yampa River to reach the Green; drift sampling in 2012 and 2014-2016 was completed for portions of the summer but few smallmouth bass were captured.

Task 4: Sample larval drift and process samples.

We collected Green River drift net samples from 15 July to 15 August (n = 81 samples) in 2016. Very few fish were captured (n = 17) but with the exception of a single white sucker larva, the remainder was native fishes. No roundtail chub or Colorado pikeminnow larvae were captured in 2016.

Task 5: Identify, count and measure preserved samples of small-bodied fish (seine hauls).

We have completed identification of 2015 samples and are progressing with 2016 samples.

Task 6: Prepare and submit annual report.

This report.

VIII. Additional noteworthy observations:

IX. Recommendations: Continue with sampling in 2016. Implement findings of the smallmouth bass synthesis report and plan a program to disadvantage smallmouth bass using flow spike(s) in the Green River.

X. Project Status: On Track and Ongoing.

XI. FY 2016 Budget Status

- A. Funds Provided: \$187,924
- B. Funds Expended: \$101,854
- C. Difference: \$86,070, will be used to complete sample work and analyses, participate in the Green River flow evaluation team, and write a spike flow study plan.
- D. Percent of the FY 2015 work completed about 50%, no additional funds needed.
- E. Recovery Program funds spent for publication charges:

XII. Status of Data Submission (Where applicable): endangered fish data submitted

XIII. Signed: Kevin R. Bestgen      14 November 2016  
Principal Investigator      Date

*(Just put name and date here, since you will be submitting the report electronically)*

*APPENDIX: E.g., more comprehensive/final project reports (NOT to be used in place of a complete annual report.). If distributed previously, simply reference the document or report.*


Table 1.—List of fishes captured in the Green River, from Browns Park downstream to Rainbow Park with electrofishing, trammel nets, and seining, 2002-2016. White crappie is new to the list. N = native, I = introduced.

| Species             | Status | Electrofishing | Trammel netting | Seining        |
|---------------------|--------|----------------|-----------------|----------------|
| Mountain whitefish  | N      | X              |                 | X              |
| Humpback chub       | N      | X              | X               |                |
| Bonytail            | N      | X              | X               | X <sup>1</sup> |
| Roundtail chub      | N      | X              | X               | X              |
| Colorado pikeminnow | N      | X              | X               | X              |
| Speckled dace       | N      | X              |                 | X              |
| Bluehead sucker     | N      | X              | X               | X              |
| Flannelmouth sucker | N      | X              | X               | X              |
| Razorback sucker    | N      | X              |                 |                |
| Mountain sucker     | N      |                |                 | X              |
| Mottled sculpin     | N      | X              |                 | X              |
| Cutthroat trout     | I      | X              |                 |                |
| Brook trout         | I      | X              |                 |                |
| Rainbow trout       | I      | X              |                 | X              |
| Brown trout         | I      | X              | X               |                |
| Northern pike       | I      | X              |                 | X              |
| Red shiner          | I      | X              |                 | X              |
| Common carp         | I      | X              | X               | X              |
| Creek chub          | I      |                |                 | X              |
| Fathead minnow      | I      |                |                 | X              |
| Sand shiner         | I      |                |                 | X              |
| Redside shiner      | I      | X              |                 | X              |
| White sucker        | I      | X              | X               | X              |
| WS x FM             |        | X              | X               |                |
| FM x BH             |        | X              |                 |                |
| WS x BH             |        | X              |                 |                |
| RZB x FM            |        | X              |                 | X              |
| Channel catfish     | I      | X              | X               | X              |
| Black bullhead      | I      | X              |                 | X              |
| Bluegill            | I      | X              |                 | X              |
| Green sunfish       | I      | X              |                 | X              |
| Smallmouth bass     | I      | X              | X               | X              |
| White crappie       | I      |                |                 | X              |
| Walleye             | I      | X              |                 |                |
| Iowa darter         | I      |                |                 | X              |


<sup>1</sup> Stocked fish.

Table 2. Number of fish of selected species captured by sampling trip (July and September) in the Lodore (LD) and Whirlpool Canyon (WH) reaches of the Green River, 2016.

| Species                        | July LD    | July WH    | Sept LD    | Sept WH    | July Total  | Sept Total  | LD Total    | WH Total   | Grand Total |
|--------------------------------|------------|------------|------------|------------|-------------|-------------|-------------|------------|-------------|
| flannemouth sucker             | 307        | 116        | 200        | 163        | 423         | 363         | 507         | 279        | 786         |
| bluehead sucker                | 99         | 125        | 51         | 75         | 224         | 126         | 150         | 200        | 350         |
| brown trout                    | 163        | 0          | 179        | 3          | 163         | 182         | 342         | 3          | 345         |
| channel catfish                | 74         | 83         | 67         | 62         | 157         | 129         | 141         | 145        | 286         |
| mountain whitefish             | 71         | 8          | 137        | 32         | 79          | 169         | 208         | 40         | 248         |
| white sucker                   | 40         | 3          | 40         | 5          | 43          | 45          | 80          | 8          | 88          |
| common carp                    | 35         | 7          | 27         | 15         | 42          | 42          | 62          | 22         | 84          |
| flannemouth x white sucker     | 24         | 0          | 37         | 4          | 24          | 41          | 61          | 4          | 65          |
| smallmouth bass                | 14         | 18         | 13         | 6          | 32          | 19          | 27          | 24         | 51          |
| mottled sculpin                | 6          | 2          | 7          | 3          | 8           | 10          | 13          | 5          | 18          |
| rainbow trout                  | 5          | 3          | 5          | 3          | 8           | 8           | 10          | 6          | 16          |
| bluehead x white sucker        | 2          | 2          | 5          | 3          | 4           | 8           | 7           | 5          | 12          |
| Colorado pikeminnow            | 3          | 0          | 2          | 1          | 3           | 3           | 5           | 1          | 6           |
| northern pike                  | 3          | 0          | 2          | 0          | 3           | 2           | 5           | 0          | 5           |
| flannemouth x bluehead sucker  | 0          | 2          | 2          | 0          | 2           | 2           | 2           | 2          | 4           |
| reidside shiner                | 0          | 0          | 3          | 0          | 0           | 3           | 3           | 0          | 3           |
| red shiner                     | 0          | 2          | 0          | 0          | 2           | 0           | 0           | 2          | 2           |
| flannemouth x razorback sucker | 0          | 1          | 0          | 0          | 1           | 0           | 0           | 1          | 1           |
| razorback sucker               | 0          | 0          | 0          | 1          | 0           | 1           | 0           | 1          | 1           |
| roundtail chub                 | 0          | 0          | 0          | 1          | 0           | 1           | 0           | 1          | 1           |
| speckled dace                  | 1          | 0          | 0          | 0          | 1           | 0           | 1           | 0          | 1           |
| walleye                        | 1          | 0          | 0          | 0          | 1           | 0           | 1           | 0          | 1           |
| <b>Totals</b>                  | <b>848</b> | <b>372</b> | <b>777</b> | <b>377</b> | <b>1220</b> | <b>1154</b> | <b>1625</b> | <b>749</b> | <b>2374</b> |




A.




B.

Figure 1. below



C.



D.

Figure 1. Green River water temperatures at the Gates of Lodore, near the Dinosaur National Monument campground, June 2002- November 2016 (Panel A, preceding page) and 2016 (panel B, preceding page). Panels C and D represent 2016 mean daily water temperature and flow at the Greendale gauge just downstream of Flaming Gorge Dam.




Figure 2.—Number of brown trout, channel catfish, common carp, northern pike, smallmouth bass, and white sucker captured per hour of raft electrofishing effort in four reaches of Lodore Canyon and two reaches of Whirlpool Canyon, Green River, Colorado and Utah, in from 1994-2016. LD is Lodore Canyon and WH is Whirlpool Canyon.




Figure 3.—Number of bluehead sucker, Colorado pikeminnow, flannelmouth sucker, mountain whitefish, and roundtail chub captured per hour of raft electrofishing effort in four reaches of Lodore Canyon and two reaches of Whirlpool Canyon, Green River, Colorado and Utah, from 1994-2016. LD is Lodore Canyon and WH is Whirlpool Canyon.






Figure 4 continued below.






Figure 4 continued below.






Figure 4 continued below.




Figure 4.—Distributions of hatching dates of Age-0 smallmouth bass estimated by otolith daily increment analysis, 2003-2011, and first hatching and hatch duration in 2012. Bass were collected from the Green River in Lodore Canyon. Left vertical axis is the frequency of fish in the histograms or water temperature (increasing but variable trace through time, red if in color); right vertical axis is Green River discharge (in cubic feet per second and is depicted by a declining or stable line time, green if in color). Data are presented and discussed in Bestgen and Hill (2016).

Appendix I. Browns Park sampling results, spring 2016.

| <b>Trammel Netting</b>                                  |              |
|---|--------------|
| <b>Species</b>  | <b>Total</b> |
| Colorado pikeminnow                                     | 6            |
|   |              |
| bluehead sucker   | 9            |
| bluehead sucker x<br>flannelmouth sucker                | 18           |
| flannelmouth sucker                                     | 135          |
|   |              |
| mountain whitefish                                      | 3            |
| brown trout   | 10           |
|   |              |
| northern pike   | 5            |
| white sucker  | 245          |
| white sucker X flannelmouth<br>sucker                   | 153          |
| white sucker X bluehead sucker                          | 11           |
| white sucker X bluehead sucker<br>X flannelmouth sucker | 1            |
| channel catfish   | 2            |
| common carp   | 6            |
| <b>Total</b>  | <b>604</b>   |
| <b>CPUE (fish/net hr)</b>                               | <b>1.71</b>  |

| <b>Boat Electrofishing</b>                              |              |
|---|--------------|
| <b>Species</b>  | <b>Total</b> |
| Colorado pikeminnow                                     | 4            |
|   |              |
| bluehead sucker   | 0            |
| bluehead sucker x<br>flannelmouth sucker                | 3            |
| flannelmouth sucker                                     | 19           |
|   |              |
| mountain whitefish                                      | 69           |
| brown trout   | 57           |
|   |              |
| northern pike   | 0            |
| white sucker  | 59           |
| white sucker X flannelmouth<br>sucker                   | 52           |
| white sucker X bluehead<br>sucker                       | 0            |
| white sucker X bluehead<br>sucker X flannelmouth sucker | 0            |
| channel catfish   | 0            |
| common carp   | 2            |
| <b>Total</b>  | <b>265</b>   |
| <b>CPUE (fish/hr)</b>                                   | <b>77.03</b> |

| <b>Trammel and Electrofishing Combined</b>              |              |
|---|--------------|
| <b>Species</b>  | <b>Total</b> |
| Colorado pikeminnow                                     | 10           |
|   |              |
| bluehead sucker   | 9            |
| bluehead sucker x<br>flannelmouth sucker                | 21           |
| flannelmouth sucker                                     | 154          |
|   |              |
| mountain whitefish                                      | 72           |
| brown trout   | 67           |
|   |              |
| northern pike   | 5            |
| white sucker  | 304          |
| white sucker X flannelmouth<br>sucker                   | 205          |
| white sucker X bluehead<br>sucker                       | 11           |
| white sucker X bluehead<br>sucker X flannelmouth sucker | 1            |
| channel catfish   | 2            |
| common carp   | 8            |
| <b>Total</b>  | <b>869</b>   |

*For projects where more than one agency/entity receives funds from the Bureau of Reclamation, append one PPR from each agency/entity. Otherwise, delete.*

## **ANNUAL PERFORMANCE PROGRESS REPORT (PPR)**

BUREAU OF RECLAMATION AGREEMENT NUMBER: R14AP00001

UPPER COLORADO RIVER RECOVERY PROGRAM PROJECT NUMBER: FR-115

Project Title: Monitoring effects of Flaming Gorge Dam releases on the Lodore and Whirlpool Canyon fish communities

Principal Investigator: Kevin R. Bestgen, K. Zelasko, C. T. Wilcox, and A. A. Hill.  
Larval Fish Laboratory  
Dept. of Fish, Wildlife, and Conservation Biology  
Colorado State University  
Ft Collins, Colorado 80523  
970-491-1848/ fax 970-491-5091  
kbestgen@colostate.edu

Project/Grant Period: Start date (Mo/Day/Yr): 1 Oct. 2014  
End date: (Mo/Day/Yr): 30 Sept. 2018  
Reporting period end date: 30 Sept. 2016  
Is this the final report? Yes \_\_\_\_\_ No X

Performance: The Larval Fish Laboratory completed three sampling trips with assistance from the USFWS, Vernal, Utah. Samples were collected and preserved and data on large-bodied fishes was collected. Samples are being identified and data are yet being analyzed, as some information was only recently collected. We also produced an annual report on activities and will prepare presentations based on data at workshops or meetings in January 2017.

## ANNUAL PERFORMANCE PROGRESS REPORT (PPR)

BUREAU OF RECLAMATION AGREEMENT NUMBER: R15PG00083

UPPER COLORADO RIVER RECOVERY PROGRAM PROJECT NUMBER: FR-115

Project Title: Monitoring effects of Flaming Gorge Dam releases on the Lodore and Whirlpool Canyon fish communities

Principal Investigator: Tildon Jones  
U.S. Fish and Wildlife Service  
1380 S 2350 W, Vernal, UT 84078  
[tildon\\_jones@fws.gov](mailto:tildon_jones@fws.gov); (435) 789-0351

Project/Grant Period:            Start date (Mo/Day/Yr): 10/01/2014  
   End date: (Mo/Day/Yr): 9/30/2019  
   Reporting period end date (Mo/Day/Yr): 09/30/2016  
   Is this the final report? Yes \_\_\_\_\_ No   X  

Performance: USFWS completed our portions of tasks 1 and 2, sampling large and small bodied fish communities. We provided personnel and equipment to successfully complete 2 electrofishing and seining passes from Lodore Canyon through Island/Rainbow Parks, as well as a trammel netting trip in Whirlpool Canyon. These tasks were completed in collaboration with CSU, who maintains data files and is ultimately responsible for submitting project reports and analyses.