

COLORADO RIVER RECOVERY PROGRAM
FY 2017 ANNUAL PROJECT REPORT

RECOVERY PROGRAM
PROJECT NUMBER: 22f

I. Project Title: INTERAGENCY STANDARDIZED MONITORING PROGRAM (ISMP)
ASSESSMENT OF ENDANGERED FISH REPRODUCTION IN RELATION TO FLAMING
GORGE OPERATIONS IN THE MIDDLE GREEN AND LOWER YAMPA RIVERS-Yampa
and middle Green River assessment of Colorado pikeminnow and razorback sucker larvae

II. Bureau of Reclamation Agreement Number(s): R14AP00001

Project/Grant Period: Start date (Mo/Day/Yr): 1 Oct. 2014
End date: (Mo/Day/Yr): 30 Sept. 2018
Reporting period end date: 30 Sept. 2017
Is this the final report? Yes No X

III. Principal Investigator(s): Kevin R. Bestgen
Larval Fish Laboratory (LFL)
Department of Fish, Wildlife, and Conservation Biology
Colorado State University
Fort Collins, CO 80523
(970) 491-1848/5295; FAX 491-5091
E-mail kbestgen@colostate.edu

Tildon Jones
USFWS
Colorado River Fishery Project
1380 S. 2350 W.
Vernal, Utah 84078
Phone: (435) 789-0354; Fax: (435) 789-4805

IV. Abstract: The goal of Flaming Gorge flow and temperature recommendations (Muth et al., 2000) that were implemented in 2006 was to improve the status and prospects for recovery of endangered fish populations in the Green River. A major emphasis of those recommendations was to enhance the reproductive and recruitment success of endangered fishes in the middle Green River, in particular razorback sucker and Colorado pikeminnow. This data will be used to assess effects of flow and temperature regimes on reproduction by razorback suckers and Colorado pikeminnow and to correlate abundance of larvae to abundance of juveniles in autumn. Larvae of razorback sucker *Xyrauchen texanus* and Colorado pikeminnow *Ptychocheilus lucius* were captured in the Green River basin in spring and summer 2017. Razorback sucker sampling was conducted with light traps primarily in the Green River between Jensen and Ouray and Colorado pikeminnow sampling was with drift nets in the lower Yampa River. Sampling was designed to provide a measure of timing of reproduction and a measure of annual reproductive success of each species. Diel variation in abundance of Colorado pikeminnow larvae in the drift was also assessed.

V. Study Schedule: Ongoing in a new agreement began in 2014, similar sampling has been conducted since 1990 for Colorado pikeminnow except for 1997, and since 1992 for razorback suckers. Anticipate continued annual sampling to build this valuable long-term monitoring dataset.

VI. Relationship to RIPRAP: Relationship to specific RIPRAP items:

Green River Action Plan: Mainstem

I. Provide and protect instream flows--habitat management.

I.A. Green River above Duchesne River.

I.A.1. Initially identify year-round flows needed for recovery while providing experimental flows.

I.A.2.a. Summer/fall flow recommendations.

I.A.3. Deliver identified flows.

I.A.3.a. Operate Flaming Gorge pursuant to the Biological Opinion to provide summer and fall flows.

I.A.3.d. Operate Flaming Gorge Dam to provide winter and spring flows and revised summer/fall flows, if necessary.

I.B. Green River below the Duchesne River.

I.B.1. Initially identify year-round flows needed for recovery while providing experimental flows.

I.B.2. State acceptance of initial flow recommendations.

I.B.2.a. Review scientific basis.

II. Restore habitat--habitat development and maintenance.

II.A. Restore and manage flooded bottomland habitat.

II.A.1. Conduct site restoration.

II.A.1.a. Old Charlie Wash.

II.A.1.a.(3) Monitor and evaluate success.

II.C. Enhance water temperatures to benefit endangered fishes.

II.C.1. Identify options to release warmer water from Flaming Gorge Reservoir to restore native fish habitat in the Green River.

V. Monitor populations and habitat and conduct research to support recovery actions-- research, monitoring, and data management.

V.A. Conduct research to acquire life history information and enhance scientific techniques required to complete recovery actions.

Green River Action Plan: Yampa and Little Snake Rivers

I. Provide and protect instream flows--habitat management.

I.D. Yampa River below Little Snake River.

I.D.1. Initially identify year-round flows needed for recovery.

I.D.2. Evaluate need for instream flow water rights.

I.D.2.a. Review scientific basis.

Green River Action Plan: Yampa and Little Snake Rivers

V.A.1. Conduct standardized monitoring.

V.B.2. Conduct appropriate studies to provide needed life history information.

VII. Accomplishment of FY 2017 Tasks and Deliverables, Discussion of Initial Findings and Shortcomings:

Project Objectives

- 1). Determine timing and duration of spawning by razorback suckers and timing of first presence (3 June) and abundance of larvae in the system as measured by capture of larvae in light traps. Additional sampling was also conducted in flood plain wetlands in early summer and autumn under this project.
- 2). Determine timing and duration of spawning by Colorado pikeminnow and timing of first presence (1 July) and abundance of larvae in the system as measured by capture of larvae downstream of spawning areas in the lower Yampa River.

Task Description (FY 2017)

- I). Collect light trap samples for razorback suckers. The CRFP office in Vernal was be responsible for this task.
- II). Collect drift net samples for Colorado pikeminnow. The Larval Fish Laboratory was responsible for this task.
- III). Identify light trap and drift net samples. Preliminary identifications will be conducted by the responsible sampling entity, with assistance from the LFL, as samples are collected to provide real-time data. Final specimen identification and curation will be conducted by the LFL.
- IV). Summarize specimen data collection in an annual report.

Accomplishments by Task.

- I). Collect light trap samples for razorback suckers. Light trap samples were collected during May and June, 2017 by the Vernal CRFP. Additional sampling near Green River flood plain wetlands was also conducted in summer 2017.
- II). Collect drift net samples for Colorado pikeminnow. Drift net samples were collected during June to August 2017 by the Larval Fish Laboratory.
- III). Identify light trap and drift net samples.

Middle Green River light trap samples, 2015. The identification of 2015 samples was nearly complete by reporting time in 2015, and was reported in final form in 2016. We retain those 2015 and 2016 data in this report for comparison to 2017 information when that is available for a longer-term report; 2017 samples have not yet been identified. The duration of the sampling season was 5 May until 12 June, 2015. The first razorback sucker larvae of the season was captured on 7 May at Cliff Creek, the earliest date ever recorded since this sampling program began in 1992. That date of first appearance was in contrast to 2014, where first larvae appearance was 28 May, a relatively average date of first appearance, and to 2011 when the first larvae were detected late on 24 June. Flows

in 2011 were extremely high and cold, which delayed reproduction by razorback suckers; flows were moderately high in 2014, and in 2015, were low in early May and warm, but frequent and heavy rains increased river levels through May and into June. Dates of first appearance of razorback sucker larvae have been successfully used to make decisions regarding Flaming Gorge Dam water management in spring for several years. Larvae captured in 2015 light trap samples ranged from 10-19 mm total length. The relatively large specimens captured on the last few sampling days (10.8-16 mm TL) indicated that spawning may finished some time before the last sampling date.

Abundance of razorback sucker larvae in 2015 was different than typical, as most larvae were captured prior to onset of high releases from Flaming Gorge Dam and increased Green River, Jensen, Utah, flows (Figure 2). One goal of those increased releases is to connect middle Green River floodplain wetlands with the river when razorback sucker larvae are produced, to aid in their growth and recruitment. Presence of larvae triggers higher releases from Flaming Gorge Dam in spring to support flows from the Yampa River, but typically higher numbers of larvae are present when flow releases peak. The relatively low production of juvenile razorback suckers in Stewart Lake wetland in 2015 could be associated with the lower abundance of larvae produced after flow levels increased.

We also prepared samples of razorback suckers captured for selenium analysis in 2014 and 2015 in conjunction with Recovery Program staff (McAbee).

Middle Green River light trap samples, 2016. The identification of 2016 samples was completed in early 2017 so we now report those results. The duration of the sampling season was mid-May until late June. First razorback sucker larvae of the season were captured on 28 May at Cliff Creek, 21 days after first larvae were collected in 2015 (7 May), and on the same date as larvae were first collected in 2014; 28 May is an average date of first collection. In contrast, first appearance of larvae in 2011 when flows were high and cold very late, was 24 June. Larvae captured in 2016 light trap samples ranged from 8-17 mm total length. Dates of first appearance of razorback sucker larvae were used to make decisions regarding Flaming Gorge Dam water management in spring since 2012.

Number of razorback sucker larvae collected was down again in 2016 from earlier peaks in 2012 and 2013 (Figure 1), which is about when large numbers of razorback suckers stocked in 2009-2010 were detected at the middle Green River razorback sucker spawning area (Webber et al. 2014, Project 169 annual report). The 2016 numbers were slightly higher than in 2015 and 2014. Accessibility to sampling sites and likely, lower fish abundance may have been factors associated with those lower numbers. In the past, larger numbers of larvae were also associated with lower water years, including 2012 and 2013, as well as in 1994, when fewer and only wild fish were present. The lower 2016 number of larvae are in contrast to the high numbers of juvenile fish produced in Stewart Lake in 2016.

We also examined the potential role that reduced stocking rates may play in the trend for recently declining numbers of razorback suckers stocked from hatcheries. Based on PIT tag detections at spawning areas, hatchery fish take 3-4 years post stocking to begin showing up at the spawning areas (Smith et al. 2016, Project 169 data). Effects of lower numbers of fish stocked from 2013-2017 may only be evident in a couple more years when mortality begins to reduce abundance of larger groups of 2009-2012 stocked fish. However, those fish have been persistent and even increasing in abundance at the spawning bar based on PIT tag detections over the 2014-2016 period (Project 169 results, see Appendix I for those data); detection rates may have also increased over time and accounted for some of the apparent increase in abundance. It is difficult to correlate numbers of spawning adults detected at the spawning areas with larvae produced and captured in this study, because many of the smaller groups of fish recently stocked (2014-2017) may not be of spawning age yet. Continued monitoring of this relationship between numbers of stocked fish and numbers of larvae produced 3-4 year later is warranted in the future.

Abundance of razorback sucker larvae in 2016 was well-timed with increased Green River, Jensen, Utah, flows (Figure 2). Presence of larvae triggers higher releases from Flaming Gorge Dam in spring to support and enhance flows from the Yampa River. One goal of those releases is to connect middle Green River floodplain wetlands with the river when razorback sucker larvae are produced, to entrain larvae into productive floodplain wetlands and aid in their growth and recruitment. The higher abundance of larvae later in the season associated with higher flows is positively related to the relatively large number of juvenile razorback suckers (>2,000) produced in Stewart Lake in autumn 2016 (Project 163 annual reports).

Middle Green River light trap samples, 2017. The identification of 2017 samples has not been completed. When those data are available we will report it, and compare with 2015 and 2016 data which is also included in this report for comparison. We note that first capture of razorback sucker was on 3 June at the Stewart Lake drain, and was from samples collected by Utah Division of Wildlife Resources as part of their Stewart Lake inundation studies (Project 165, those larvae were identified by B. Haines, USFWS, Vernal, Utah). First razorback suckers captured in Cliff Creek, in this study, were 6 June 2017. Sampling extended into July on Project 22f.

Lower Yampa River drift net sampling, 2015. Samples were collected in the Yampa River about 0.2 to 0.8 km upstream from the Green River (n = 246 total samples collected in 2015), the same site that samples were collected from in 1990 to 1996 (Bestgen et al. 1998) and in 1998 to 2014. Sampling commenced on 17 June and extended through 16 August. The first Colorado pikeminnow larva was collected on 27 June, a relatively average date for first capture of larvae and not unexpected given the warm (but high levels) water temperatures early in summer.

Abundance of Colorado pikeminnow larvae was about average in 2015, but less than that for 2013 and 2014 (Figure 3). Yampa River flows were moderate to low early in the reproductive season. We plotted capture dates of larvae as a function of time and

overlaid that with Green River flow. Green River flows at Jensen, Utah, were relatively low in early July and near the level (about 3000 cfs) when habitat conditions are suitable for good survival of young Colorado pikeminnow (Bestgen and Hill 2016). Presence of early lower flows may have increased survival of those larvae, because a large year-class of YOY pikeminnow in autumn was documented (Project 138 results). A flow spike in the Yampa River on 10 July 2015 may have reduced survival of pikeminnow larvae, similar to that for smallmouth bass upstream (Annual report, Project 140), because few pikeminnow larvae were captured from about 10-22 July. However, a bimodal peak of reproduction is not uncommon (see 2016 and 2017 data) so low abundance of larvae could also be due to lack of reproduction in the preceding days. About 50% or more of Green River, Jensen flows in July were from the Green River, and the remainder was from the Yampa River, the exception being the 10 July flow spike.

Abundance of YOY Colorado pikeminnow in 2015 in the middle Green River, Utah, ISMP samples collected in autumn was very high, especially when compared to recent years (M. Breen personal communication, Project 138 annual report). Above average abundance of larvae does not appear to be one of the reasons that YOY were so abundant, as similar levels of larvae were produced in 2016 and few YOY resulted (see below). One possible reason for increased YOY production in 2015 is that summer flows in the Green River, especially those early in July when pikeminnow were hatching, were moderate to low. This is unlike 2016 and especially 2017 when early July flows were high (see below). Backwaters were also present and abundant at that time in 2015, and lower flow levels likely allowed for colonization by the invertebrate food base that Colorado pikeminnow require, which likely also assisted with production of a large YOY pikeminnow year class. This situation is in contrast to 2016 and 2017, years with similar levels of larvae production but had higher or very high flows in early July, respectively. The YOY year classes in autumn essentially failed in those years in the Green River.

Lower Yampa River drift net sampling, 2016. Samples were collected in the Yampa River about 0.2 to 0.8 km upstream from the Green River (n = 233 total samples collected in 2016), the same site that samples were collected from in 1990 to 1996 (Bestgen et al. 1998) and in 1998 to 2015. Sampling commenced on 23 June and extended through 16 August. The first Colorado pikeminnow larva was collected on 30 June, a relatively average date for first capture of larvae and not unexpected given the warm (but high levels) water temperatures early in summer.

Abundance of Colorado pikeminnow larvae was about average in 2016, similar to that in 2015, but less than that for 2013 and 2014 (Figure 3). High Yampa River flows were present early in the reproductive season. We plotted capture dates of larvae as a function of time and overlaid that with Green River flow. Green River flows at Jensen, Utah, were high in early July and well above the flow levels when abundant backwater habitat is available. Even though flows were lower in the later portion of July, reduced or absent backwater habitat in early July likely reduced survival of those early larvae. About 50% of those early July flows were produced in the Yampa River, and the other half derived from releases from Flaming Gorge Dam.

Abundance of YOY Colorado pikeminnow in 2016 in the middle Green River, Utah, ISMP samples collected in autumn was very low (M. Breen personal communication, Project 138 annual report). Abundance of larvae does not appear to be one of the reasons for reduced YOY abundance as larvae were at average or higher than average abundances in 2016. One possible reason for reduced YOY production is that early July flows were high, during a period when about 50% of Colorado pikeminnow larvae were produced. Thus, backwaters may not have been available during that time. Further, newly formed backwater habitat is likely food-poor because invertebrates have not colonized those areas and established populations, which would logically reduce survival of larvae. Flow patterns, especially timing of flows when larvae are first present, as well as magnitude, may be very important for production of good year classes of YOY pikeminnow.

Reasons for lack of YOY Colorado pikeminnow produced from larvae available in the later portion of 2016 are not known. Sampling in autumn 2016 indicated backwaters were arrayed in a different geomorphic pattern than they normally are, and some that were historically present were gone (Matt Breen, personal communication, Project 138 annual report). Habitat shifts may thus, be an additional reason for poor production of YOY Colorado pikeminnow in 2016.

Most sampling data collected in this program (1990-2012, none in 1997) were recently incorporated into a report entitled "Reproduction, abundance, and recruitment dynamics of young Colorado pikeminnow in the Green River Basin, Utah and Colorado, 1979-2012. That report (Bestgen and Hill 2016) has been reviewed by the Biology Committee of the Upper Colorado River Endangered Fish Recovery program, comments incorporated into a final report, which was approved and finalized.

Lower Yampa River drift net sampling, 2017. Samples were collected in the Yampa River about 0.2 to 0.8 km upstream from the Green River (n = 243 total samples collected in 2017), the same site that samples were collected from in 1990 to 1996 (Bestgen et al. 1998) and in 1998 to 2016. Sampling commenced on 21 June and extended through 14 August. The first Colorado pikeminnow larva was collected on 1 July, a relatively average date for first capture of larvae and not unexpected given the relatively low water level and higher temperatures early in summer in the Yampa River. Flows were high and temperatures were low in the adjacent Green River well after pikeminnow larvae were first present due to extended and high releases from Flaming Gorge Dam.

Abundance of Colorado pikeminnow larvae was below average in 2017, and lower than was observed in 2016 or 2015 (Figure 3). Higher Yampa River flows were present early in the reproductive season but dropped relatively quickly. We plotted capture dates of pikeminnow larvae as a function of time and overlaid that with Green River flow. Green River flows at Jensen, Utah, were high in early July and likely higher than when any backwater habitat is available. This was also coincident with the time of highest production of larvae, as 80% of all larvae produced in 2017 were present before base flows declined. Even after that, base flows remained relatively high through the summer. Even though flows were lower in the later portion of July, reduced or absent backwater


habitat in early July likely reduced survival of those early larvae both in the middle and lower Green River. This was borne out by low numbers of YOY Colorado pikeminnow captured in autumn in the middle (n = 1) or lower (n < 25) Green River. This was not unlike 2016 as well, when abundance of YOY Colorado pikeminnow in the middle Green River, Utah, ISMP samples was very low (M. Breen personal communication, Project 138 annual report). One possible reason for reduced YOY production is that early July flows were high and backwaters were not available during that time. Further, newly formed backwater habitat is likely food-poor because invertebrates have not colonized those areas and established populations. Thus, suitable backwater habitat may not have been available for the few larvae that were available until later in 2017, if at all. Flow pattern timing, essentially reducing flows to adequate levels, as well as appropriate magnitude, may be very important for production of good year classes of YOY pikeminnow. A comparison of flows in 2015 to those in 2016 and 2017 shows starkly different patterns because 2015 flows were low early and resulted in a large year class, whereas 2016 and 2017 flows were high later, and resulted in essentially a year-class failure in the Green River. The reduction in abundance of larvae produced in the Yampa River and captured in 2017 is also worrisome.

- VIII. Additional noteworthy observations: Four razorback sucker larvae (8-18 mm TL) were captured in drift nets at the Yampa River drift net sampling site in 2016, a gravel bar formerly known to support reproduction by razorback suckers. Capture dates of larvae ranged from 30 June to 12 July. A few adult razorback suckers are detected there annually with portable PIT tag detectors (Smith et al. 2015; 2016, Project 168). Numerous chub larvae, presumably roundtail chub *Gila robusta*, were also noted in samples in 2016. The 2017 data have not yet been summarized for species other than Colorado pikeminnow but will be updated when available.
- IX. Recommendations: Continue sampling as planned in 2018. Continue to integrate this work into recruitment patterns noted for juveniles per Bestgen and Hill (2016).
- X. Project Status: Ongoing and on-track.
- XI. FY 2017 Budget Status
- A. Funds Provided: \$ 128,747
 - B. Funds Expended: \$ 96,560
 - C. Difference: \$32,187 remaining funds for sample analysis that remains.
 - D. Percent of the FY 2017 work completed, and projected costs to complete: About 65% complete.
 - E. Recovery Program funds spent for publication charges: None.
- XII. Status of Data Submission: Data were made available to investigators.
- XIII. Signed: Kevin R. Bestgen 12 November 2017
Principal Investigator Date

APPENDIX: Major recent products based on these data include:

- Bestgen, K. R., and A. A. Hill. 2016. Reproduction, abundance, and recruitment dynamics of young Colorado pikeminnow in the Green River Basin, Utah and Colorado, 1979-2012. Final report to the Upper Colorado River Endangered Fish Recovery Program, Project FW BW-Synth, Denver, CO. Department of Fish, Wildlife, and Conservation Biology, Colorado State University, Fort Collins. Larval Fish Laboratory Contribution 183.
- Bestgen, K. R. 2015. Aspects of the Yampa River flow regime essential for maintenance of native fishes. Final Report submitted to the National Park Service, Fort Collins, CO, The Nature Conservancy, Boulder, CO, and Western Resource Advocates, Boulder, CO. Department of Fish, Wildlife, and Conservation Biology, Colorado State University, Fort Collins. Larval Fish Laboratory Contribution 181.
- Bestgen, K. R., K. A. Zelasko, and G. C. White. 2012. Monitoring reproduction, recruitment, and population status of razorback suckers in the Upper Colorado River Basin. Final Report to the Upper Colorado River Endangered Fish Recovery Program, U. S. Fish and Wildlife Service, Denver. Larval Fish Laboratory Contribution 170.
- Bestgen, K. R., G. B. Haines, and A. A. Hill. 2011. Synthesis of flood plain wetland information: Timing of razorback sucker reproduction in the Green River, Utah, related to stream flow, water temperature, and flood plain wetland availability. Final report to the Recovery Implementation Program for Endangered Fishes in the Upper Colorado River Basin. U. S. Fish and Wildlife Service, Denver, CO. Larval Fish Laboratory Contribution 163.

A.



B.

Green River razorback sucker stocking






Figure 1. Panel A depicts number of razorback sucker larvae captured from 1993 to 2016 in the middle Green River, Utah, in light traps (all fish including those of questionable taxonomic identity included; 2017 sample identification is not finished so is not included here). Panel B depicts number of razorback suckers stocked each year throughout the Green River (about ½ in each of the middle and lower Green River each year), 1995-2017.

A.



B.



C.




Figure 2. Seasonal distribution of razorback sucker larvae captured in 2015 (upper panel, a), 2016 (middle panel, b), and 2017 (lower panel, first capture only as indicated by arrow) in middle Green River, Utah, in light trap samples (all fish including those identified as “razorback sucker?” were included). All samples from all locations were combined for each day. Gaps in captures after the first capture of razorback sucker larvae may indicate lack of sampling rather than absence of the species. Green River flows at Jensen, Utah (solid line) are plotted, as are releases from Flaming Gorge Dam (Greendale, Utah gauge, dashed line), with the difference in the two lines being mainly flows of the Yampa River.






Figure 3. Number of Colorado pikeminnow larvae captured from 1990 to 2017 (no sampling in 1997, includes specimens from all diel samples, 2017 sample identification is complete) in the lower Yampa River, Colorado, during summer in drift nets.

a.



b.



c.




Figure 4. Seasonal distribution of Colorado pikeminnow larvae captured in 2015 (panel a), 2016 (panel b), and 2017 (panel c) drift net samples from the lower Yampa River, Colorado. Sampling for 2015 began on 17 June and ended 16 August. Sampling for 2016 began on 23 June and ended 16 August. Sampling for 2017 began on 21 June and ended 14 August. Several of the largest capture dates were diel sampling occasions (12 samples total each day). Flows of the Green River at Jensen (solid line) and Greendale (dashed line) are also depicted, with the difference in the two flows being the contribution of the Yampa River.

FY 2017 Ann. Rpt. Project # 22f, pikeminnow and razorback larvae sampling - 16
 Appendix I. Detections of razorback suckers at Razorback Bar complex, middle Green River, Utah, from 2014-2016, showing abundance of various year-classes of fish stocked over time. Razorback suckers may take about 3 or more years post stocking to mature, based on first relatively high abundances of fish detected at spawning areas three years after stocking.




Figure 1. Year of stocking for razorback sucker detected with the PIT antennas in 2014.




Figure 3. Year of stocking for razorback sucker detected at Razorback Bar PIT tag antennas in 2015.




Figure 3. Year of stocking for razorback sucker detected at Razorback Bar PIT tag-antennas in 2016.

FY 2017 Ann. Rpt. Project # 22f, pikeminnow and razorback larvae sampling - 17
ANNUAL PERFORMANCE PROGRESS REPORT (PPR)

BUREAU OF RECLAMATION AGREEMENT NUMBER: R14AP00001

UPPER COLORADO RIVER RECOVERY PROGRAM PROJECT NUMBER: 22f

Project Title: INTERAGENCY STANDARDIZED MONITORING PROGRAM (ISMP)
ASSESSMENT OF ENDANGERED FISH REPRODUCTION IN RELATION
TO FLAMING GORGE OPERATIONS IN THE MIDDLE GREEN AND
LOWER YAMPA RIVERS.

Principal Investigator: Kevin R. Bestgen
Larval Fish Laboratory (LFL)
Department of Fish, Wildlife, and Conservation Biology
Colorado State University
Fort Collins, CO 80523
(970) 491-1848/5295; FAX 491-5091
E-mail kbestgen@colostate.edu

Project/Grant Period: Start date (Mo/Day/Yr): 1 Oct. 2014
End date: (Mo/Day/Yr): 30 Sept. 2018
Reporting period end date: 30 Sept. 2017
Is this the final report? Yes _____ No X

Performance: The goal of this project is to document timing and intensity of reproduction by razorback suckers and Colorado pikeminnow in the lower Yampa and middle green rivers. In 2017, we collected 243 drift net samples and additional seine samples in the lower Yampa River, and began identification of those. The 2016 results are reported here as well. In addition, we identified samples collected by the U. S. Fish and Wildlife Service in 2016 and began sample identification for 2017 samples. Another objective of this project is to summarize data collections. This was done in the attached report.

FY 2017 Ann. Rpt. Project # 22f, pikeminnow and razorback larvae sampling - 18
ANNUAL PERFORMANCE PROGRESS REPORT

BUREAU OF RECLAMATION AGREEMENT NUMBER: R15PG00083

UPPER COLORADO RIVER RECOVERY PROGRAM PROJECT NUMBER: 22f

Project Title: Light trap and drift net sampling for razorback sucker and Colorado pikeminnow larvae

Principal Investigator:

M. Tildon Jones
1380 S 2350 W, Vernal, UT 84078
tildon_jones@fws.gov; 435-789-0351 x14

Project/Grant Period:

Start date: 10/01/2014

End date: 09/30/2019

Reporting period end date: 09/30/2017

Is this the final report? Yes _____ No X

Performance:

US Fish & Wildlife Service completed our portion of Task 1, collect light trap and seine samples from the Green River and its wetlands. We began sampling May 26, 2017 and continued light trapping both the river and floodplain sites of interest through July 12. Data analysis and reporting will be completed by CSU-LFL.